RECEIVED CENTRAL FAX CENTER

APR 1 8 2007

7

Docket No. C&R-117 Serial No. 10/600,790

In the Claims

- 1 (currently amended). A An isolated polypeptide, which polypeptide that:
- (i) comprises or consists of the amino acid sequence as recited in SEQ ID NO:2;
- (ii) is a fragment of SEQ ID NO:2-thereof that is an interferon gamma like-secreted protein of the four-helical bundle cytokine fold, or having an antigenic determinant in common withpolypoptides of (i); or
- (iii) is a functional equivalent of (i) or (ii) has greater than 90% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof;

wherein said isolated polypeptide or fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

2-5 (canceled).

6 (currently amended). The fragment or functional equivalent of claim 4 isolated polypeptide according claim 1, wherein said polypeptide has greater than 95% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavatin A or phytohemagglutinin stimulated peripheral blood mononuclear cells there is greater than 95% sequence identity.

7 (currently amended). The fragment or functional equivalent of claim 4 isolated polypeptide according claim 1, wherein said polypeptide has greater than 98% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-a, IL-2, IL-4, IFN-y, aspartate

J:\C&R\117\Amd-Resp\Amd.doc/DNB/gyl

8

Docket No. C&R-117 Serial No. 10/600,790

aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-y secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells, wherein there is greater than 98% sequence identity.

8 (currently amended). The fragment or functional equivalent of claim-4 isolated polypeptide according to claim 1, wherein said polypeptide has greater than 99% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells wherein there is greater than 99% sequence identity.

9-50 (canceled).

- 51 (new). The isolated polypeptide according to claim 1, wherein said polypeptide comprises the amino acid sequence as recited in SEQ ID NO: 2.
- 52 (new). The isolated polypeptide according to claim 1, wherein said polypeptide consists of the amino acid sequence as recited in SEQ ID NO: 2.
- 53 (new). The isolated polypeptide according to claim 1, wherein said polypeptide is a fragment of SEQ ID NO: 2 and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.
- 54 (new). The isolated polypeptide according to claim 1, wherein said polypeptide is a fragment of a polypeptide having greater than 90% sequence identity to SEQ ID NO: 2 and said fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels

J:\C&R\117\Amd-Resp\Amd.doc/DNB/gyl

9

Docket No. C&R-117 Serial No. 10/600,790

of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

- 55 (new). A composition comprising a pharmaceutical carrier and a polypeptide that:
 - (i) comprises or consists of the amino acid sequence as recited in SEQ ID NO:2;
 - (ii) is a fragment of SEQ ID NO:2; or
 - (iii) has greater than 90% sequence identity with the polypeptide of SEQ ID NO:

2 or a fragment thereof;

wherein said isolated polypeptide or fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

56 (new). The composition according to claim 55, wherein said polypeptide has greater than 95% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

57 (new). The composition according to claim 55, wherein said polypeptide has greater than 98% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

10

Docket No. C&R-117 Serial No. 10/600,790

- 58 (new). The composition according to claim 55, wherein said polypeptide has greater than 99% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.
- 59 (new). The composition according to claim 55, wherein said polypeptide comprises the amino acid sequence as recited in SEQ ID NO: 2.
- 60 (new). The composition according to claim 55, wherein said polypeptide consists of the amino acid sequence as recited in SEQ ID NO: 2.
- 61 (new). The composition according to claim 55, wherein said polypeptide is a fragment of SEQ ID NO: 2 and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.
- 62 (new). The composition according to claim 55, wherein said polypeptide is a fragment of a polypeptide having greater than 90% sequence identity to SEQ ID NO: 2 and said fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF-α, IL-2, IL-4, IFN-γ, aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) in vivo, or the ability to induce IFN-γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.